

## HOW TO USE MISOPROSTOL FOR A SELF-ABORTION BEFORE 12 WEEKS OF PREGNANCY




1 hour before (recommended for pain):2 pills of 400mg of Ibuprofen

4 tablets of 200mcg of Misoprostol



Under the tongue Dissolution in 30mn



**Every 3 hours until expulsion** Maximum 3 doses



To increase the chances of a complete abortion, it is recommended that you take a second dose of misoprostol, even if you have already started bleeding or if something has been expelled after the first dose, especially if the pregnancy is less than 9 weeks.


## HOW TO USE MISOPROSTOL FOR A SELF-ABORTION AFTER 12 WEEKS OF PREGNANCY






1 hour before (recommended for pain):2 pills of 400mg of Ibuprofen

2 tablets of 200mcg of Misoprostol



Under the tongue Dissolution in 30mn






**Every 3 hours until expulsion** Maximum 3 doses




To increase the chances of a complete abortion, it is recommended that you take a second dose of misoprostol, even if you have already started bleeding or if something has been expelled after the first dose, especially if the pregnancy is less than 9 weeks.

## HOW TO USE MIFEPRISTONE AND MISOPROSTOL FOR A SELF-ABORTION BEFORE 12 WEEKS OF PREGNANCY



1 tablet of mifepristone (200mg)






2 pills of 400mg of Ibuprofen (recommended for pain)

One hour after:

4 tablets of 200mcg of Misoprostol




Under the tongue Dissolution in 30mn

**Every 3 hours until expulsion - maximum 3 dosess** 

To increase the chances of a complete abortion, it is recommended that you take a second dose of misoprostol, even if you have already started bleeding or if something has been expelled after the first dose, especially if the pregnancy is less than 9 weeks.

# HOW TO USE MIFEPRISTONE AND MISOPROSTOL FOR A SELF-ABORTION AFTER 12 WEEKS OF PREGNANCY.


1



1 tablet of mifepristone (200mg)



2




2 pills of 400mg of Ibuprofen (recommended for pain)

**One hour after:** 


3

2 tablets of 200mcg of Misoprostol



Under the tongue Dissolution in 30mn

**Every 3 hours until expulsion - maximum 3 dosess** 





To increase the chances of a complete abortion, it is recommended that you take a second dose of misoprostol, even if you have already started bleeding or if something has been expelled after the first dose, especially if the pregnancy is less than 9 weeks.

## WARNING SIGNS AND HOW TO REACT

### GO TO THE NEAREST HEALTH CENTER OR HOSPITAL IF:



#### **Heavy vaginal bleeding**

- bleeding that soaks two sanitary pads or cloth pads for normal periods in one hour, and if it repeats in the following hour
- bleeding accompanied by dizziness or lightheadednes




Greenish or yellowish foul-smelling discharge from the vagina



Blood clots larger than an orange



Intense and persistent abdominal pain that worsens despite taking painkillers



A fever above 38°C that begins within 24 hours of taking misoprostol

If you do not wish to say that you have taken medication to terminate a pregnancy, you can simply say that you are bleeding. Medical staff cannot tell the difference between a natural miscarriage and a medical abortion.